

Lower bound for electron spin entanglement from beamsplitter current correlations

Guido Burkard¹ and Daniel Loss²

¹IBM T. J. Watson Research Center, P. O. Box 218, Yorktown Heights, NY 10598

²Department of Physics and Astronomy, University of Basel,
Klingelbergstrasse 82, CH-4056 Basel, Switzerland

We determine a lower bound for the entanglement of pairs of electron spins injected into a mesoscopic conductor. The bound can be expressed in terms of experimentally accessible quantities, the zero-frequency current correlators (shot noise power or cross-correlators) after transmission through an electronic beam splitter. The effect of spin relaxation (T_1 processes) and decoherence (T_2 processes) during the ballistic coherent transmission of the carriers in the wires is taken into account within Bloch theory. The presence of a variable inhomogeneous magnetic field allows the determination of a useful lower bound for the entanglement of arbitrary entangled states. The decrease in entanglement due to thermally mixed states is studied. Both the entanglement of the output of a source (entangler) and the relaxation (T_1) and decoherence (T_2) times can be determined.

Quantum nonlocality has been an intriguing issue since the early days of quantum mechanics [1]. Nonlocal effects can come into play when a quantum system is composed of at least two subsystems which are spatially separated. Despite their simplicity, the Bell states of two distant quantum two-state subsystems (A and B)

$$|\Psi_{\pm}\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle \pm |\downarrow\uparrow\rangle), \quad (1)$$

$$|\Phi_{\pm}\rangle = \frac{1}{\sqrt{2}}(|\uparrow\uparrow\rangle \pm |\downarrow\downarrow\rangle), \quad (2)$$

exhibit the essential phenomenology of quantum nonlocality (e.g., they violate Bell's inequalities [2]) thus providing an ideal testing ground for quantum nonlocality. Here, we represent the two-state systems as spins $1/2$ with basis states “spin up” $|\uparrow\rangle$ and “spin down” $|\downarrow\rangle$ with respect to an arbitrary fixed direction in space.

With the development of quantum information theory [3], and in particular with quantum communication, it has become clear that EPR pairs can also play the role of a *resource* for operations that are impossible with purely classical means. In this context, two-state systems are referred to as quantum bits (qubits), and quantum nonlocality is related to the concept of entanglement (defined below). A number of quantum information processes—quantum teleportation [4], quantum key distribution [5], quantum dense coding [6], etc.—have been successfully implemented using pairs of photons with entangled polarizations, i.e., in states such as Eqs. (1) and (2). Photons have the advantage that they are easily moved from one place to another, allowing for experiments involving space-like separations between detection events [2].

More recently, there has been increasing interest in the use of the spin of electrons in a solid-state environment for spin-based electronics [7] and as qubits for quantum computing [8]. Subsequently, quantum communication on a mesoscopic scale, typically on the order of micrometers in semiconductor structures (e.g. quantum wires), was proposed [9]. Rather than achieving space-like separation between detection events on the two sites (this




FIG. 1: Inset: Proposed setup with two-electron scattering at a beamsplitter (BS) with transmittivity T . Electrons are injected pairwise from the entangler (E) into contacts 1 and 2 of the BS. The mean current $I = \langle I_{\alpha} \rangle$ and one of the current correlators $S_{\alpha\beta}$ are measured at the outgoing contacts $\alpha, \beta = 3, 4$. Plot: Entanglement of formation E of the electron spins versus singlet fidelity F and the reduced correlator $f = S_{33}/2eIT(1 - T)$. The curve illustrates the relation between noise and entanglement for Werner states. For general states, the curve represents a lower bound for the entanglement, i.e. allowed values for E and f (or, equivalently, F) are represented by points in the shaded region. Measuring f determines a lower bound for the entanglement E .

would require sub-picosecond detection), the idea here is to use quantum entanglement between parts of a coherently operating solid-state device (in the most extreme case, a quantum computer). It is then relevant to study the transport of spin-entangled electrons in a many-electron system and possible means of entanglement detection. Two-particle interference at a beamsplitter (BS) combined with the measurement of current fluctuations [10] (in general, the full counting statistics [11]) was identified as a detector for entanglement.

In this paper, we go one step further, providing a *lower bound* for the *amount* E of spin entanglement carried by individual pairs of electrons, related to the zero-frequency current correlators when measured in a BS setup (Fig. 1,

Inset) by injecting the electrons separately into the two ingoing leads (1 and 2) and measuring either the current autocorrelator $S_{\alpha\alpha}$ in one of the outgoing leads ($\alpha = 3, 4$) or the cross-correlator S_{34} . It is assumed that the size of the scattering region is smaller than both the coherence length and the mean free path, allowing for ballistic and coherent transport. In the following, T will denote the transmittivity of the BS, i.e. the probability to be scattered from lead 1 to lead 4 (or from 2 to 3). The ideal BS for the proposed setup does not give rise to backscattering (e.g. from lead 1 back into lead 1, or from 1 into 2, etc.). We will also analyze the effect of such backscattering processes, as they give rise to background shot noise which is unrelated to entanglement. During their transport, the electron spins will be exposed to decoherence and relaxation due to spin-dependent scattering caused by magnetic impurities, nuclear spins, or the spin-orbit coupling (see [12] for a review). We include these effects within a Bloch equation formalism [13]. Comparison between our theory and experiment will (i) test proposed entanglers [14, 15, 16, 17, 18, 19, 20] and (ii) determine spin relaxation (T_1) and decoherence (T_2) times.

The materials and structures required for testing our theory, although at the forefront of current capabilities, appear to be feasible. The largest efforts seem to be necessary to realize the electron spin entangler [10] for which there exists a number of theoretical ideas, using normal-[14, 15] or carbon-nanotube-superconductor junctions [16, 17, 18], or single [19], or coupled quantum dots [10, 20]. The electronic BS and the measurement of BS current correlators have been experimentally demonstrated in a GaAs/AlGaAs heterostructure [21]. Coherent transport of electron spins over more than 100 μm in GaAs has been observed [22].

Traditionally, current correlations, and in particular the quantum partition (shot) noise have been used to gain information about a scatterer beyond its conductance [23]. Here, we use a known scatterer (the BS) to gain information about the quantum state (more precisely, its entanglement) of the scattered particles. The correlation function between the currents $I_\alpha(t)$ and $I_\beta(t)$ in two leads $\alpha, \beta = 1, \dots, 4$ of the BS is defined as

$$S_{\alpha\beta}(\omega) = \lim_{\tau \rightarrow \infty} \frac{\hbar\nu}{\tau} \int_0^\tau dt e^{i\omega t} \text{Re Tr} [\delta I_\alpha(t) \delta I_\beta(0) \chi], \quad (3)$$

where $\delta I_\alpha = I_\alpha - \langle I_\alpha \rangle$, $\langle I_\alpha \rangle = \text{Tr}(I_\alpha \chi)$, ν is the density of states in the leads, and χ is the density matrix of the injected electron pair (below, we suppress the orbital part of χ , see [10] for Coulomb effects). Writing χ in the Bell basis, Eqs. (1) and (2), $\chi = F|\Psi_-\rangle\langle\Psi_-| + G_0|\Psi_+\rangle\langle\Psi_+| + \sum_{i=\pm} G_i|\Phi_i\rangle\langle\Phi_i|$, and $S_{\alpha\beta} \equiv S_{\alpha\beta}(\omega = 0)$, we arrive at

$$S_{\alpha\beta} = FS_{\alpha\beta}^{|\Psi_-\rangle} + G_0S_{\alpha\beta}^{|\Psi_+\rangle} + \sum_{i=\pm} G_iS_{\alpha\beta}^{|\Phi_i\rangle}, \quad (4)$$

$$S_{\alpha\beta}^{|\Psi\rangle} \equiv \lim_{\tau \rightarrow \infty} \frac{\hbar\nu}{\tau} \int_0^\tau dt \text{Re} \langle \Psi | \delta I_\alpha(t) \delta I_\beta(0) | \Psi \rangle. \quad (5)$$

Using the standard scattering approach [23], we have found earlier [10] that the singlet state $|\Psi_-\rangle$ gives rise to enhanced shot noise (and cross-correlators) at zero temperature, $S_{33}^{|\Psi_-\rangle} = -S_{34}^{|\Psi_-\rangle} = 2eIT(1-T)f$, with the *reduced correlator* $f = 2$, as compared to the “classical” Poissonian value $f = 1$ [24]. The average currents are given by $I = \langle I_3 \rangle = \langle I_4 \rangle = e/h\nu$. We also know that all triplet states are noiseless, $S_{\alpha\beta}^{|\Psi_+\rangle} = S_{\alpha\beta}^{|\Phi_\pm\rangle} = 0$ ($\alpha, \beta = 3, 4$). Both the current autocorrelations (shot noise) and cross-correlations are only due to the singlet component of the incident two-particle wavefunction,

$$S_{33} = -S_{34} = FS^{|\Psi_-\rangle} = 2eIT(1-T)f, \quad f = 2F. \quad (6)$$

Including backscattering with probability R_B , we find

$$S_{33} = 2eI[2F(1-R_B)T(1-T) + R_B/2], \quad (7)$$

$$S_{34} = -2eI2F(1-R_B)T(1-T), \quad (8)$$

where $I = (e/h\nu)(1-R_B)$. Since $f' = S_{34}/2eIT(1-T) = 2F(1-R_B) \leq f$ is smaller than f without backscattering and the entanglement of formation E is a monotonic function of f (see below and Fig. 1), we still obtain a lower bound on E (the bound will become less informative as R_B increases). Note that this does not hold for the autocorrelator S_{33} . However, one can determine R_B , e.g. by measuring the shot noise power using normal Fermi lead inputs [21] and then obtain f from either S_{33} or S_{34} .

The entanglement of a bipartite state $\chi \in \mathcal{H}_A \otimes \mathcal{H}_B$ can be quantified by its entanglement of formation [25] $E(\chi) = \min_{\{(|\chi_i\rangle, p_i)\} \in \mathcal{E}(\chi)} \sum_i p_i S_N(|\chi_i\rangle)$, where $\mathcal{E}(\chi)$ denotes the set of ensembles $\{(|\chi_i\rangle, p_i)\}$ for which $\chi = \sum_i p_i |\chi_i\rangle\langle\chi_i|$. We have used the von Neumann entropy of the reduced density matrix $\rho_B = \text{Tr}_A |\psi\rangle\langle\psi|$, $S_N(|\psi\rangle) = -\text{Tr}_B \rho_B \log \rho_B$ (logarithms are in base 2). A state with $E > 0$ ($E = 1$) is (maximally) entangled, whereas a state with $E = 0$ is separable (in the case of a pure state, it is a product $\psi_A \otimes \psi_B$). The Bell states Eq. (1) and (2) are maximally entangled. Neither local operations nor classical communication (LOCC) between subsystems A and B can increase E . In quantum information theory, $E(\chi)$ is the maximal ratio N/M of the number N of EPR pairs (maximally entangled states) required to form M copies of χ as $N \rightarrow \infty$; E is the quantity that measures *how much* of the resource (quantum entanglement) is available.

For arbitrary χ , $E(\chi)$ cannot be expressed as a function of only its singlet fidelity $F = \langle \Psi_- | \chi | \Psi_- \rangle$. However, this is possible for the so-called Werner states [26]

$$\rho_F = F|\Psi_-\rangle\langle\Psi_-| + \frac{1-F}{3} \left(|\Psi_+\rangle\langle\Psi_+| + \sum_{i=\pm} |\Phi_i\rangle\langle\Phi_i| \right), \quad (9)$$

being the unique rotationally invariant states with singlet fidelity F . It is known [25] that $E(F) \equiv E(\rho_F) = H_2(1/2 + \sqrt{F(1-F)})$ if $1/2 < F \leq 1$ and $E(F) \equiv$




FIG. 2: Homogeneous magnetic field $\delta h = 0$ and $\tilde{P} = 1$. (a) f of the spin singlet state $|\Psi_{-}\rangle$ after ballistic transmission through a BS as a function of the spin decoherence time T_2 , in units of the ballistic transmission time $t_0 = L/v_F$. Different curves correspond to different values of the spin relaxation time T_1 (same units). Note that we only plotted the curves for $T_2 \leq 2T_1$. (b) f for a spin triplet state $|\Psi_{+}\rangle$. Since $f \leq 1$, the lower bound on entanglement is zero, i.e. we cannot learn anything about entanglement of injected triplets at $\delta h = 0$. (c) Lower bound on the entanglement of formation E .

$E(\rho_F) = 0$ if $0 \leq F < 1/2$, with the dyadic Shannon entropy $H_2(x) = -x \log x - (1-x) \log(1-x)$. Together with Eq. (6), this enable us to express the entanglement of ρ_F in terms of the reduced correlator f (Fig. 1).

We generalize this result to arbitrary mixed states χ of two spins (qubits). Any state χ can be transformed into the Werner state ρ_F with the same singlet fidelity F by a random bipartite rotation [25, 27], i.e. by choosing

a random $U \in SU(2)$ and applying $U \otimes U$ to χ . Since this operation involves only LOCC, E cannot increase,

$$E(F) \leq E(\chi). \quad (10)$$

The entanglement of formation $E(F)$ of the corresponding Werner state therefore provides a *lower bound* on the entanglement of χ (Fig.1). Thus, a noise signal exceeding the Poissonian limit ($f > 1$) in the BS setup can in principle be interpreted as a sign of entanglement between the electron spins injected into leads 1 and 2 [30].

We now include relaxation and decoherence into our analysis. At time $t = 0$, we start with a spin singlet (upper sign) or triplet (lower sign) state

$$\chi(0) = |\Psi_{\mp}\rangle\langle\Psi_{\mp}|. \quad (11)$$

We describe the dynamics of $\chi(t)$ in a field $\mathbf{B} \parallel \hat{z}$ and in the presence of spin decoherence (T_2 processes) and relaxation (T_1) phenomenologically within a single-spin Bloch equation for the polarization $\mathbf{P} = (\langle\sigma_x\rangle, \langle\sigma_y\rangle, \langle\sigma_z\rangle)$,

$$\dot{\mathbf{P}} = \mathbf{P} \times \mathbf{h} - R(\mathbf{P} - \tilde{\mathbf{P}}) \equiv -\Omega(\mathbf{P} - \tilde{\mathbf{P}}), \quad (12)$$

with $\langle\sigma_i\rangle = \text{Tr}(\sigma_i \rho)$, $\mathbf{h} = g\mu_B \mathbf{B} = (0, 0, h)$, the stationary polarization $\tilde{\mathbf{P}} = (0, 0, \tilde{P})$ (note that $\tilde{\mathbf{P}} \times \mathbf{h} = 0$), and the relaxation matrix [31] $R_{ij} = \delta_{ij} R_i$ with $R_1 = R_2 = T_2^{-1}$ and $R_3 = T_1^{-1}$. Solving Eq. (12), we obtain

$$\mathbf{P}(t) = e^{-\Omega t} \mathbf{P}(0) + (1 - e^{-\Omega t}) \tilde{\mathbf{P}}, \quad (13)$$

or, in terms of the spin density matrix,

$$\rho(t) = (P_0 + \mathbf{P}(t) \cdot \boldsymbol{\sigma})/2 \equiv \Lambda_h(t)[\rho(0)], \quad (14)$$

with the superoperator ($a(t) = 1 - e^{-t/T_1}$) [32]

$$\Lambda_h(t)[\rho] = \begin{pmatrix} \frac{1}{2}(\rho_{\uparrow\uparrow} + \rho_{\downarrow\downarrow})(1 + a(t)\tilde{P}) + \frac{1}{2}(\rho_{\uparrow\uparrow} - \rho_{\downarrow\downarrow})e^{-t/T_1} & e^{-t/T_2 + iht}\rho_{\uparrow\downarrow} \\ e^{-t/T_2 - iht}\rho_{\downarrow\uparrow} & \frac{1}{2}(\rho_{\uparrow\uparrow} + \rho_{\downarrow\downarrow})(1 - a(t)\tilde{P}) - \frac{1}{2}(\rho_{\uparrow\uparrow} - \rho_{\downarrow\downarrow})e^{-t/T_1} \end{pmatrix}, \quad (15)$$

with the matrix elements $\rho_{ij} = \langle i|\rho|j\rangle$ and $(i, j = \uparrow, \downarrow)$. We apply $\Lambda_h(t)$ to both spins individually,

$$\chi(t) = (\Lambda_{h_1}(t) \otimes \Lambda_{h_2}(t)) [\chi(0)], \quad (16)$$

where h_i is the field at electron i . Using Eq. (6) and $F(t) = \langle\Psi_{-}|\chi(t)|\Psi_{-}\rangle$ at time $t \geq 0$, we obtain

$$f(t) = \pm e^{-2t/T_2} \cos(\delta h t) + \frac{1}{2}(1 + e^{-2t/T_1}) - \frac{1}{2}(1 - e^{-t/T_1})^2 \tilde{P}^2, \quad (17)$$

where $\delta h = h_1 - h_2$. If the decoherence time $T_2^{(1,2)}$ of the

two electrons is different, then T_2 in Eq. (17) becomes $T_2^{\text{EPR}} = (1/T_2^{(1)} + 1/T_2^{(2)})^{-1}$. We define T_1^{EPR} similarly if $\tilde{P} = 0$. However, if $\tilde{P} = 1$, then $\exp(-t/T_1)$ is replaced by $\exp(-t/T_1^{(1)}) + \exp(-t/T_1^{(2)})$.

A homogeneous field, $\delta h = 0$, does not affect f . For slow relaxation, $T_1 \gg t$, we find $f(t) = 1 \pm e^{-2t/T_2}$. In Fig. 2a we plot f for $\delta h = 0$ and $\tilde{P} = 1$ versus T_2 in units of the ballistic transmission time [24] $t_0 = L/v_F$ (L =length of ballistic trajectory, v_F =Fermi velocity). For unentangled triplet states, $\chi(0) = |\uparrow\uparrow\rangle\langle\uparrow\uparrow|, |\downarrow\downarrow\rangle\langle\downarrow\downarrow|$, we find $f \leq 1/2$ for all T_1, T_2 , and $\tilde{P}$ (Fig. 2b).




FIG. 3: (a) Reduced current correlator f versus field inhomogeneity $\delta h = h_1 - h_2$ (in units of $\hbar/t_0 g \mu_B$, t_0 =ballistic transmission time, g =g-factor, μ_B =Bohr magneton) for an injected singlet state and $\tilde{P} = 1$. Solid lines represent $T_1 \gg 1$ and $T_2 = 1, 3, 10, \infty$, grey dashed lines $T_1 = 3$ and $T_2 = 1, 3$. For injected triplets, the plot is phase-shifted by π , providing tight lower bounds at $\delta h = \pi$. Tight lower bounds for *any* input state can be determined by varying the direction of $\delta \mathbf{h}$. (b) Plot of f for a thermally mixed initial state versus T_2 (in units of t_0) for $T_1 \gg t$ and $\tilde{P} = 1$. The various curves correspond to $k_B T/J = 0, 0.25, 0.33, 0.5, 1$, where J and T denote the exchange energy and temperature during the preparation of the state. Inset: The maximal f (at $T_1, T_2 \gg t_0$) versus $k_B T/J$. There is no entanglement ($f \leq 1$, $E = 0$) above the critical temperature $T_c = 0.91 J/k_B$.

An inhomogeneous field $\delta \mathbf{h} \neq 0$ (or, equivalently, a local controllable Rashba spin-orbit coupling [28]) has the effect of continuously rotating singlets into triplets and vice versa (Fig. 3). This a lower bound of the triplet entanglement, $\chi(0) = |\Psi_+\rangle\langle\Psi_+|$, which is as tight as Eq. (10) for the singlet,

$$E \geq \max_{\delta \mathbf{h}} E(f(\delta \mathbf{h})/2), \quad (18)$$

where $f(\delta \mathbf{h})$ is the measured noise power (or cross-correlator), $E(F)$ is the entanglement of the Werner state ρ_F (Fig. 1). If a field inhomogeneity $\delta \mathbf{h}$ can be created pointing in *arbitrary* directions in space, then the above result represents a tight lower bound for *any* injected entangled state. In particular, each maximally entangled state $|\Psi\rangle$ will be detected in this way, since there exists a $u = \exp(-i\delta \mathbf{h} \cdot \boldsymbol{\sigma}) \in SU(2)$ such that $u^\dagger \otimes u |\Psi\rangle = |\Psi_- \rangle$. This rotation can also be done unilaterally, i.e. there is a $v \in SU(2)$ with $v \otimes 1 |\Psi\rangle = |\Psi_- \rangle$ (see also [28]).

Finally, we study the case where the spin state of the injected pair of carriers Eq. (11) is mixed, because it is prepared at a temperature T comparable to the energy splitting between spin states, typically (if the Zeeman effect is negligible) the exchange energy J , i.e. the singlet-triplet splitting. In this case, $\chi(0) = \rho_F$ with $F = (1 + 3e^{-J/k_B T})^{-1}$ where k_B is Boltzmann's constant. We only show the resulting f for $T_1 \gg t$ here (the full expression will be reported elsewhere [29]),

$$f(t) = \frac{1 + e^{-2t/T_2} + e^{-J/k_B T} (1 - e^{-2t/T_2})}{1 + 3e^{-J/k_B T}}, \quad (19)$$

which is the statistical mixture of Eq. (17) for the singlet and triplet with the appropriate Boltzmann weights

(Fig. 3b). Above the critical temperature $T_c = 0.91 J/k_B$ there is no entanglement even for $T_1, T_2 \rightarrow \infty$.

Acknowledgments: We thank D. P. DiVincenzo and B. M. Terhal for valuable discussions. DL acknowledges funding from the Swiss NSF, NCCR Nanoscience, and DARPA QUIST and SPINS.

-
- [1] A. Einstein, B. Podolski, and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [2] J. S. Bell, Rev. Mod. Phys. **38**, 447 (1966); A. Aspect, J. Dalibard, G. Roger, Phys. Rev. Lett. **49**, 1804 (1982).
 - [3] C.H. Bennett, D.P. DiVincenzo, Nature **404**, 247 (2000).
 - [4] C. H. Bennett *et al.*, Phys. Rev. Lett. **70**, 1895 (1993); D. Boumeester *et al.*, Nature **390**, 575 (1997); D. Boschi *et al.*, Phys. Rev. Lett. **80**, 1121 (1998).
 - [5] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
 - [6] K. Mattle *et al.*, Phys. Rev. Lett. **76**, 4656 (1996).
 - [7] *Semiconductor Spintronics and Quantum Computation*, eds. D. D. Awschalom, D. Loss, and N. Samarth (Springer, Berlin, 2002).
 - [8] D. Loss, D. P. DiVincenzo, Phys. Rev. A **57**, 120 (1998).
 - [9] D. V. DiVincenzo and D. Loss, J. Mag. Magn. Matl. **200**, 202 (1999) [cond-mat/9901137].
 - [10] G. Burkard, D. Loss, and E. V. Sukhorukov, Phys. Rev. B **61**, R16303 (2000) [cond-mat/9906071].
 - [11] F. Taddei and R. Fazio, Phys. Rev. B **65**, 075317 (2002).
 - [12] M. E. Flatté, J. M. Byers, W. H. Lau, Chap. 4 in Ref. [7].
 - [13] C. P. Slichter, *Principles of Nuclear Magnetic Resonance*, 3rd ed. (Springer, Berlin, 1990).
 - [14] P. Recher, E. V. Sukhorukov, and D. Loss, Phys. Rev. B **63**, 165314 (2001).
 - [15] G. B. Lesovik, T. Martin, and G. Blatter, Eur. Phys. J. B **24**, 287 (2001).
 - [16] P. Recher and D. Loss, Phys. Rev. B **65**, 165327 (2002).
 - [17] C. Bena *et al.*, Phys. Rev. Lett. **89**, 037901 (2002).
 - [18] V. Bouchiat *et al.*, cond-mat/0206005.
 - [19] W. D. Oliver, F. Yamaguchi, and Y. Yamamoto, Phys. Rev. Lett. **88**, 037901 (2002).
 - [20] D. Saraga and D. Loss, cond-mat/0205553.
 - [21] R. C. Liu *et al.*, Nature (London), **391**, 263 (1998).
 - [22] J. M. Kikkawa and D. D. Awschalom, Nature **397**, 139 (1999).
 - [23] Ya. M. Blanter, M. Büttiker, Phys. Rep. **336**, 1 (2000).
 - [24] For a discussion of the range of validity of this result and an estimate of the ballistic transfer time t_0 , see Section 9 in J. C. Egues *et al.*, cond-mat/0210498.
 - [25] C. H. Bennett *et al.*, Phys. Rev. A **54**, 3824 (1996).
 - [26] R. F. Werner, Phys. Rev. A **40**, 4277 (1989).
 - [27] C. H. Bennett *et al.*, Phys. Rev. Lett. **76**, 722 (1996).
 - [28] J. C. Egues, G. Burkard, and D. Loss, Phys. Rev. Lett. **89**, 176401 (2002).
 - [29] G. Burkard and D. Loss, unpublished.
 - [30] If the current is carried by quasiparticles with charge e^* , f will be renormalized by a factor e/e^* .
 - [31] Within standard weak-coupling theory, $T_2 = [(2T_1)^{-1} + T_{\parallel}^{-1}]^{-1} \leq 2T_1$ ($T_{\parallel}^{-1}$ is the longitudinal decoherence rate).
 - [32] The superoperator $\Lambda_h(t)$ is linear and trace-preserving on density matrices with arbitrary trace since we have not imposed the trace condition $P_0 = 1$ at this point.